

Finite size scaling of entanglement entropy at the Anderson transition with interactions

Rui-Lin Chu, An Zhao and Shun-Qing Shen

Department of Physics, The University of Hong Kong, Pokfulam Road, Hong Kong, China

(Dated: October 10, 2012)

We study the entanglement entropy (EE) of disordered one-dimensional spinless fermions with attractive interactions. With intensive numerical calculation of the EE using the density matrix renormalization group method, we find clear signatures of the transition between the localized and delocalized phase. In the delocalized phase, the fluctuations of the EE becomes minimum and independent of the system size. Meanwhile the EE's logarithmic scaling behavior is found to recover to that of a clean system. We present a general scheme of finite size scaling of the EE at the critical regime of the Anderson transition, from which we extract the critical parameters of the transition with good accuracy, including the critical exponent, critical point and a power-law divergent localization length.

PACS numbers: 71.30.+h, 72.15.Rn, 03.65.Ud

Although the Anderson transitions between localized and delocalized phases for single particles has been well understood during the past few decades, the case of disordered interacting particles remains incompletely understood and is much more challenging.[1, 2] For the simplest model with interaction - one dimensional(1D) spinless fermions, it is predicted that repulsive interaction enhances localization while attractive interaction decreases localization. For sufficiently strong attractive interactions, a transition from localization to delocalization was predicted from the analytic renormalization group approach.[3, 4] However, the many-body nature of this system makes numerical verification of these predictions a challenging task. Previous numerical studies utilizing phase sensitivity of the ground state energy to boundary conditions has succeeded to map out this delocalized phase.[5, 6] By combining the density matrix renormalization group(DMRG) with the transfer matrix method, Carter and MacKinnon were able to extract the localization length and obtain the phase diagram in the disorder-interaction phase space.[7]

The development of quantum information theory in the past decade has shed new light on this problem.[8] Entanglement entropy(EE) has been found able to characterize both quantum criticality and topological phases in a handful of quantum many-body systems.[9–14] Recently there has been increasing interest of characterizing the disordered quantum systems through the EE.[15–19] The EE is a measure of the quantum correlation in a system. For a pure state $|\Psi\rangle$ of bipartite system AB, the EE (von Neumann entropy) is defined as

$$S = -\text{Tr} \rho_A \log \rho_A = -\text{Tr} \rho_B \log \rho_B, \quad (1)$$

where $\rho_{A(B)} = \text{Tr}_{B(A)} |\Psi\rangle\langle\Psi|$ is the reduced density matrix of subsystem A(B). For a region of length L_A of a 1D metal of length L with open boundary condition, the

EE grows logarithmically as

$$S_A(L_A, L) = \frac{c}{6} \log[\sin(\frac{\pi L_A}{L})] + \tilde{c}, \quad (2)$$

where $\tilde{c}$ is a non-universal constant and c is a universal constant given by the central charge of the associated conformal field theory.

In a recent paper,[19] Berkovits proposes a formulation that extracts the localization length(LL) of a localized 1D system from the saturation behavior of S_A . However, this method is best suitable when the system shows strong localization, i.e. the LL $\xi \ll L$. When ξ becomes comparable with L , the EE suffers from strong finite size effect. Moreover, there will be no observable saturation against L_A in S_A when ξ exceeds L . In fact if there's a transition from localized into delocalized phase, ξ will become divergent. When $\xi \gg L$, the system has become metallic and its EE should behave just as that of a clean 1D metal.

The main result of this Letter is a numerical finite size scaling of the EE in the critical regime of Anderson transition in one dimension. Consider the EE between two halves of a 1D system of length L , in the localized phase where $\xi \ll L$, it is $S \sim (c/6) \log \xi$; in the delocalized phase where $\xi \gg L$, it is $S \sim (c/6) \log L$. Then in the critical regime between these two phases, we need to find a universal scaling function that takes into account the finite size effect and is able to bridge both phases. By universal we mean this function should fit to all system sizes. By fitting this scaling function to the EE of different system sizes, we demonstrate that it is possible to extract the critical parameters of the Anderson transition.

We start by describing our model Hamiltonian, which is a 1D spinless fermion wire of length L with attractive

nearest neighbor(NN) interactions

$$\mathcal{H} = -t \sum_{i=1}^{L-1} (c_i^\dagger c_{i+1} + h.c.) + \sum_{i=1}^L \epsilon_i c_i^\dagger c_i + U \sum_{i=1}^{L-1} (n_i - 1/2)(n_{i+1} - 1/2) - \mu \sum_{i=1}^L c_i^\dagger c_i, \quad (3)$$

where c and $c^\dagger$ are the annihilation and creation operators, and $n_i = c_i^\dagger c_i$. The first two terms constitute the standard Anderson model while the third term represents the attractive NN interaction ($U < 0$). The disorder is introduced as random on-site potentials ϵ_i with a uniform distribution in the range $[-W/2, W/2]$. We set the hopping amplitude $t = 1$. The last term represents the chemical potential which controls the particle density of the system. Disorder has no net contribution to the chemical potential since we tune $\sum_{i=1}^L \epsilon_i = 0$. Because the total number of particles conserves in this system, we will restrict ourselves to the fixed total particle number at half-filling and zero temperature.

In the clean limit, up to a Jordan-Wigner transformation, this model is equivalent to the $S = 1/2$ XXZ spin-chain model, for which there exists the Bethe-ansatz solution that can be used to guide our numerical study. For $|U| < 2$, the system is gapless (i.e. metallic); For $U > 2$, the charge-density-wave (CDW) phase renders the system a Mott insulator; For $U < -2$, the ground state becomes unstable to phase separation, being either empty or full filling.[20] For the disordered case, an insulator to metal transition is predicted for $U \approx -1$.[3, 4] A second metal to insulator transition happens at even stronger interaction, where the fermions start to form clusters and become localized again. In this Letter, our focus is in the range $-2 < U < 0$.

The EE is numerically obtained through the density matrix renormalization group (DMRG) algorithm, which allows the calculation of ground state and relevant physical quantities for large interacting systems with high accuracy.[21] In our case, we consider open wires of variant sizes up to $L = 1024$. During the calculation, we retain 250 (for $L < 1024$) and 330 (for $L = 1024$) states for each block, and perform lattice sweeping until both the ground state energy and the EE between the two halves of the system (i.e. $S_A(L/2, L)$) are converged. The EE data for disordered systems is averaged over 400 random disorder configurations.

For the clean case, S_A is described by Eq. 2 with central charge $c = 1$. For the disordered case, we can base on the results of previous studies for an estimate of the phase boundary between localized and delocalized phase.[5, 7] We start by fixing $W = 1$ and gradually enhance the interaction. Clear saturation behavior of S_A is found in the weak interaction regime. As is shown in Fig. 1 where the solid lines represent the clean case and the filled dots represent the disordered case. As suggested by




FIG. 1: (color online) The average EE as a function of L_A for system size $L=512$, the error bar shows the fluctuation of EE. The dash lines represent the fitting curve in Eq. 4. $W = 1.0$ (a) In the localized phase $U = -0.2$, the EE shows early saturation and large fluctuations; (b) In the delocalized phase $U = -1.2$, the EE behaves almost like clean systems with minimum fluctuations; The inset shows the fitted parameter ζ from Eq. 4, the dash line is for the clean case and the dots are for the disordered case.

Berkovits,[19] the saturation behavior can be described by

$$S'_A(L_A, L) = s(\xi) \log[\text{erf}(\frac{L_A}{\zeta(\xi, L)})] + \tilde{c}', \quad (4)$$

where $\text{erf}(x)$ is the error function and ζ can be viewed as the effective delocalized scale of the system. s is a fitting parameter and $\tilde{c}'$ is a non-universal constant. It is noted that this formula also fit the clean case, with $\zeta \propto L$ and $s \approx 1/6$. [19] In the strongly localized regime ($\xi \ll L$), $\zeta \approx \xi$, which is the case of weak U regime. But when U is gradually strengthened, ζ quickly rises, indicating an increase in ξ . When exceeding $U \approx -0.8$, ζ becomes indistinguishable with that of the clean case (see inset of Fig. 1), which means the saturation in the EE no longer exists (Fig. 1b). In this case, the LL ξ has exceeded the system size but ζ now contains no further information about it. In fact, S'_A in this case can be fitted by Eq. 2 (with $c \approx 1$) better than Eq. 4 deviating from the clean case only by a small constant, which means that both the central charge and the EE's logarithmic scaling behavior is the same with the clean case in the delocalized phase. This is in agreement with our expectation that a delocalized system, although strongly disordered, should

still has its EE resemble that of a clean system.




FIG. 2: (color online). The EE S_H of both clean (solid lines) and corresponding disordered (dot lines) systems with variant system sizes. The inset shows the fluctuation value of S_H for the disordered case. $W = 1.0$.

An interesting observation is that the fluctuation in the EE is rather large when $-1 < U < 0$ but almost vanishes when $-1.5 < U < -1$, as is indicated by the error bars in Fig. 1. We will show in the following that the delocalized phase is accompanied by a minimum fluctuation in the EE, (Fig. 2 inset) which can be viewed as a signature of Anderson transition in this system.

Although based on the fitting using Eq. 4 we can see a clear increase in the LL when $\xi < L$, it would require calculating the EE of an infinite system in order to extract a divergent LL when approaching the critical point. We thus need alternative numerical approaches to avoid this. Since the LL ξ corresponds to the correlation length in the localized regime,[19] a generalization of finite size scaling (FSS) theory for EE is possible near the critical point according to Calabrese and Cardy.[10] We use $S_A(L/2, L)$ as a scaling quantity, which is denoted as S_H . In the clean case, S_H has the form

$$S_H(L) = \frac{1}{6} \log L + c_1(U), \quad (5)$$

where c_1 is a constant depending on U (see Fig. 2 solid lines). And in the disordered case it has the form

$$S'_H(L, x) = S_H(L) + f(x) + c_2(U, W), \quad (6)$$

where $x = \xi^{-1}L/2$. The scaling function $f(x)$ satisfies that when $x \rightarrow 0$, $f(0) = 0$ and when x is large, $f(x) \approx -(1/6) \log x$, such that both the clean limit and infinite system length limit (where $S_H \sim (1/6) \log \xi$) are recovered. In theory one can obtain universal series expansion for $f(x)$ in the small and large x limit.[10] More




FIG. 3: (color online). ΔS_H as a function of W when fixing $U = -1.2$ for different system sizes. When $W < 1.5$, ΔS_H becomes size independent, i.e., in this regime $f(x)$ is vanishing.

importantly, this universality should cover both the non-interaction and interaction case. c_2 is a correction term depending on W (see Fig. 3) and weakly depending on U , clearly $c_2(U, 0) = 0$. Based on our data, we can show that when fixing W and change U , c_2 can be treated as a small constant in the critical regime.

In Fig. 2, we plot S_H for both clean and disordered systems when the interaction strength U is varied. Scaling sizes of the system are taken. In both the weak and strong U regime, $S_H(L, \xi)$ deviates strongly from $S_H(L)$ for all system sizes. But in the regime roughly when $-1.5 < U < 1.0$, this strong deviation disappears and instead the two differ only by a small constant (≈ 0.02) that is system size independent. From Eq. 6 we can see this is only possible when $f(x)$ is vanishing and ξ is infinite, and thus we can see it as a signature of the delocalized phase. Another observation is the fluctuation of the EE becomes minimum in this range, and interestingly, is also independent of the system size (see inset of Fig. 2).[17] The delocalized phase boundary that is signaled by $f(x) = 0$ can be mapped out by fixing U and sweeping W as is shown in Fig. 3. By doing this we can also see how c_2 grows with W .

We now seek to do a FSS analysis for the EE data we show in Fig. 2. The central task for FSS is to find the scaling function $f(x)$ that will fit to all system sizes. Since in the critical regime of the Anderson transition $\xi \gg L$, we need to find a good ansatz of $f(x)$ for the small x case. Our strategy is to use the non-interaction case for searching the form of $f(x)$ and then apply $f(x)$ in the fitting of the interaction case. For the non-interaction case the LL is given by $\xi \approx 105/W^2$ [23] and the EE can be more easily calculated through the formalism devel-




FIG. 4: (color online). Data collapse for the FSS of the EE of the non-interaction case for small x case. The solid curve is a best fit with $f(x) = \alpha x^\beta$ and the fitted parameters: $\alpha = -0.16$, $\beta = 0.79$. The inset shows the data collapse for large range of x .

oped by Peschel[24]. And then $f(x)$ can be obtained by fitting $\Delta S_H = S'_H - S_H$ as a function of x . The result is presented in Fig. 4, where each EE data is averaged over 2000 random disorder samples and W is varied in the range $0.1 \sim 2$. The data from different system sizes collapse nicely to a universal function that is in the logarithmic form. We find the function

$$f(x) = a \log(1 + \alpha x^\beta) \quad (7)$$

can fit well to the data in the small and intermediate x case with $c_2(0, W) = 0$, in the large x case the data fits well to $f(x) = -(1/6) \log x + \text{const.}$ For the small x case ($0 < x < 2$), a simple form $f(x) = \alpha x^\beta$ can already fit well.

For the interaction case Eq. 7 also fits well to the data (see Fig. 5). Due to the system's instability to phase separation at large interaction strength, which is exaggerated by disorder, we find it impossible to stick to half-filling when $U < -1.8$. And thus we demonstrate the FSS at the critical regime of the first Anderson transition where the EE data is sufficient ($-1.3 \leq U \leq -0.4$). At the delocalized side of U_c , we assume $\xi = \infty$, and at the localized side of U_c , we assume a power-law relation

$$\xi = K|U - U_c|^{-\nu}. \quad (8)$$

Then we can surface fit $f(x) = \alpha x^\beta$ to the EE as a function of (U, L) directly utilizing the data of all system sizes. The result is presented in Fig. 5. The fitting is rather stable and the fitted curve coincides well with the data for all system sizes. Together with the data collapse shown in the inset of Fig. 5, this confirms the validity




FIG. 5: (color online). The fitting of FSS (solid lines) function to the EE of the interaction case. The filled dots corresponds to that of Fig. 2 respectively. The fitting parameters are: $U_c = -1.12$, $\nu = 2.16$, $\alpha = -0.16$, $\beta = 0.81$, $c_2 = -0.02$, $K = 143$. The inset shows the data collapse to the scaling function $f(x) = \alpha x^\beta$.

of the scaling function $f(x)$ and our assumption for ξ . The fitted critical point $U_c = -1.12$ agrees well with the analytical result and previous numerical results.[3, 5, 7] The fitting also gives a critical exponent $\nu = 2.16$. With these parameters, we can calculate ξ with Eq. 8 at $U = 0$ and find it agrees with the analytical result $\xi = 105/W^2$. According to Calabrese and Cardy, $f(x)$ is a universal function for small x [10], this is consistent with our fitting. For both the non-interaction and interaction case, the parameter α and β agrees well in both cases.

In summary, we have demonstrated that the EE is an efficient quantity in characterizing the Anderson transition in systems with interactions. The Anderson transition has clear signatures in the EE. The first is the disappearance of saturation and regain of logarithmic scaling with unaffected central charge (when going insulator to metal); The second is size-independent minimum fluctuation and minimum deviation from the clean system's EE in the metal phase; The third, which is the main result of this Letter, is the existence of FSS of the EE from which the critical parameters (such as critical exponent, critical point and LL) can be extracted with good accuracy.

We gratefully acknowledge discussions with Cenke Xu. This work is supported by Research Grant Council of Hong Kong under Grant No. HKU 705110P.

[1] P. Anderson, Phys. Rev. 109, 1492 (1958).

[2] P. A. Lee and T. V. Ramakrishnan, Rev. Mod. Phys. 57,

- 287 (1985).
- [3] W. Apel and T. M. Rice, Phys. Rev. B, 26, 7063 (1982).
 - [4] T. Giamarchi and H. J. Schulz, Phys. Rev. B 37, 325 (1988).
 - [5] P. Schmitteckert, T. Schulze, C. Schuster, P. Schwab, and U. Eckern, Phys. Rev. Lett., 80, 560 (1998); P. Schmitteckert, R. A. Jalabert, D. Weinmann, and J. L. Pichard, Phys. Rev. Lett. 81, 2308 (1998).
 - [6] C. Schuster, R. A. Römer, and M. Schreiber, Phys. Rev. B 65, 115114 (2002).
 - [7] J. M. Carter and A. MacKinnon, Phys. Rev. B, 72, 024208 (2005).
 - [8] For a review see: L. Amico, R. Fazio, A. Osterloh, and V. Vedral, Rev. Mod. Phys. 80, 517 (2008).
 - [9] G. Vidal, J. I. Latorre, E. Rico, and A. Kitaev, Phys. Rev. Lett, 90, 227902 (2003).
 - [10] P. Calabrese and J. Cardy, J. Stat. Mech. (2004) P06002.
 - [11] S. Ryu and Y. Hatsugai, Phys. Rev. B 73, 245115 (2006).
 - [12] G. Refael and J. E. Moore, Phys. Rev. Lett., 93, 260602 (2004).
 - [13] F. Pollmann and J. E. Moore, New J. Phys. 12, 025006. (2010).
 - [14] A. M. Turner, F. Pollman and E. Berg, Phys. Rev. B 83, 075102 (2011).
 - [15] M. Fagotti, P. Calabrese¹, and J. E. Moore, Phys. Rev. B 83, 045110 (2011).
 - [16] F. Igloi and Y. C. Lin, J. Stat. Mech. (2008) P06004.
 - [17] Similar results is also reported in previous study of EE at critical points in disordered systems without interactions: X. Jia, A. R. Subramaniam, I. A. Gruzberg, and S. Chakravarty, Phys. Rev. B 77, 014208 (2008).
 - [18] J. H. Bardarson, F. Pollmann, and J. E. Moore Phys. Rev. Lett., 109, 017202 (2012).
 - [19] R. Berkovits, Phys. Rev. Lett., 108, 176803 (2012).
 - [20] C. N. Yang and C. P. Yang, Phys. Rev. 150, 321; Phys. Rev. 150, 327 (1966).
 - [21] S. R. White, Phys. Rev. Lett. 69, 2863 (1992); S. R. White and R. M. Noack, Phys. Rev. Lett. 68, 3487 (1992); S. R. White, Phys. Rev. B 48, 10 345 (1993).
 - [22] F. Pollmann, S. Mukerjee, A. M. Turner, and J. E. Moore, Phys. Rev. Lett., 102, 255701 (2009).
 - [23] R. A. Römer and M. Schreiber, Phys. Rev. Lett. 78, 515 (1997).
 - [24] I. Peschel, J. Phys. A: Math. Gen. 36 L205 (2003).